

# Noether's Theorem for *SMOOTH*, DISCRETE and Finite Element Models

Elizabeth Mansfield



## Acknowledgements:

Arieh Iserles, Cambridge

Reinout Quispel, LaTrobe

Students, Morningside Centre

for Mathematics, Beijing.

Peter Hydon, Surrey

Leverhulme Trust

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>03 JAN 2005</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Noether's Theorem for Smooth, Discrete and Finite Element Models</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University Of Kent</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM001749, Lie Group Methods And Control Theory Workshop Held on 28 June 2004 - 1 July 2004., The original document contains color images.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>31</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

# Noether's Theorem

links SYMMETRIES and conservation laws  
for Euler Lagrange Systems.

What is a conservation law?

Answer: a divergence expression which is  
zero on solutions of the system.

The heat equation

$$u_t + (-u_x)_x = 0$$

is its own conservation law. Integrating,

$$\frac{\partial}{\partial t} \int_{\Omega} u + (-u_x)_{\partial\Omega} = 0$$

where we assume  $u$  is sufficiently nice that we  
can interchange  $\partial_t$  and  $\int$ , and we have  
applied Stokes' Theorem. In words:

Rate of change                      Net of comings and goings  
of total heat in  $\Omega$                       =                      across the boundary  
no sources or sinks

# The usual examples

## Symmetry

leaves  $Ldx$  invariant

## Conserved Quantity

the quantity behind

the  $\frac{D}{Dt}$

in the Divergence

$$\left\{ \begin{array}{l} t^* = t + c \\ \text{translation in time} \end{array} \right.$$

Energy

$$\left\{ \begin{array}{l} x_i^* = x_i + c \\ \text{translation in space} \end{array} \right.$$

Linear Momenta

$$\left\{ \begin{array}{l} \mathbf{x}^* = \mathcal{R}\mathbf{x} \\ \text{rotation in space} \end{array} \right.$$

Angular Momenta

$$\left\{ \begin{array}{l} a^* = \phi(a, b) \\ b^* = \psi(a, b) \\ \phi_a \psi_b - \phi_b \psi_a \equiv 1 \\ \text{Particle relabelling} \end{array} \right.$$

Potential vorticity

# Variational Complexes 1-2-3 !

are locally exact

*SMOOTH* cf. P.J. Oliver, Applications . . .

$$\begin{array}{ccccccc}
 \xrightarrow{\text{Curl}} & \Lambda^2 & \xrightarrow{\text{Div}} & \Lambda^3 & \xrightarrow{\hat{d}} & \hat{\Lambda}_1 & \xrightarrow{\hat{d}} & \hat{\Lambda}_2 & \xrightarrow{\hat{d}} \\
 & & & & & \downarrow \pi & & \downarrow \pi & \\
 & & & & & \Lambda^1_* & \xrightarrow{\delta} & \Lambda^2_* & \xrightarrow{\delta}
 \end{array}$$

*DISCRETE* Hydon and ELM, J. FoCM

$$\begin{array}{ccccccc}
 \xrightarrow{\Delta} & \mathbf{Ex}^2 & \xrightarrow{\Delta} & \mathbf{Ex}^3 & \xrightarrow{\hat{d}} & \hat{\Lambda}_1 & \xrightarrow{\hat{d}} & \hat{\Lambda}_2 & \xrightarrow{\hat{d}} \\
 & & & & & \downarrow \pi & & \downarrow \pi & \\
 & & & & & \Lambda^1_* & \xrightarrow{\delta} & \Lambda^2_* & \xrightarrow{\delta}
 \end{array}$$

*Finite Element* ELM and GRW Quispel, CRM Proc.

$$\begin{array}{ccccccc}
 \xrightarrow{d} & \tilde{\mathcal{F}}^2 & \xrightarrow{d} & \tilde{\mathcal{F}}^3 & \xrightarrow{\hat{d} \circ \int} & \widehat{\mathcal{F}}_1 & \xrightarrow{\hat{d}} & \widehat{\mathcal{F}}_2 & \xrightarrow{\hat{d}} \\
 & & & & & \downarrow \pi & & \downarrow \pi & \\
 & & & & & \mathcal{F}^1_* & \xrightarrow{\delta} & \mathcal{F}^2_* & \xrightarrow{\delta}
 \end{array}$$

Exactness can be used to find conservation laws for non Euler-Lagrange systems via clever ansatze!

cf Hereman, Sanders, Sayers and Wang, *CRM Proceedings*; Hydon *J. Phys. A*

Exactness is proved by the use of so-called homotopy operators  $H_i$ ,

$$\begin{array}{ccccccc} & & \text{Div} & & E & & \\ & \rightarrow & \Lambda^2 & \rightarrow & \Lambda^3 & \rightarrow & \Lambda_1^* \\ \leftarrow & & \leftarrow & & \leftarrow & & \leftarrow \\ & & H_1 & & H_0 & & \end{array}$$

which satisfy

$$(\text{Div}H_1 + H_0E)\omega = \omega, \quad \text{all } \omega \in \Lambda^3$$

Thus if  $E(\omega) = 0$ , then  $\omega = \text{Div}(H_1(\omega))$ .

Idea: solve  $E(\text{clever ansatz}) = 0$  for parameters and arbitrary functions. Then you have a conservation law using  $H_1$ .

More on  $\hat{d}$  and  $\pi$

*SMOOTH*

$$\begin{aligned}
 \hat{d}(Ldx) &= \hat{d}\left(\frac{1}{2}(u_x^2 + u_{xx}^2)dx\right) \\
 &= (u_x du_x + u_{xx} du_{xx})dx \\
 &= (-u_{xx} du + u_{xxxx} du)dx \\
 &\quad + \frac{D}{Dx} (u_x du - 2u_{xx} du_x + \frac{D}{Dx} (u_{xx} du)) \\
 &= E(L)dudx + \frac{D}{Dx}\eta_L
 \end{aligned}$$

General Formula, explicit, exact, symbolic, for  $\eta_L$  known.

$E = \pi \circ \hat{d}$ , where  $\pi$  projects out the divergence term.

More than one dependent variable

$$\begin{aligned}
 \hat{d}L(x, u, v, \dots)dx &= E^u(L)dudx E^v(L)dvdx \\
 &\quad + \frac{D}{Dx}\eta_L
 \end{aligned}$$

More on  $\hat{d}$  and  $\pi$

DISCRETE

$$\begin{aligned}\hat{d}(Ldx) &= \hat{d}\left(\frac{1}{2}u_n^2 + u_n u_{n+1}\right) \Delta_n \\ &= (u_n du_n + u_{n+2} du_n + u_n du_{n+2}) \Delta_n \\ &= (u_n + u_{n+2} + u_{n-2}) du_n \Delta_n \\ &\quad + (S - \text{id})(\dots) \\ &= E(L_n) du_n \Delta_n + \Delta(\eta_{L_n})\end{aligned}$$

General Formula, explicit, exact, symbolic, for  $\eta_{L_n}$  known.

$E = \pi \circ \hat{d}$ , where  $\pi$  projects out the total difference term.

More than one dependent variable

$$\begin{aligned}\hat{d}(L_n \Delta_n) &= E^u(L_n) du_n \Delta_n + E^v(L_n) dv_n \Delta_n \\ &\quad + \Delta(\eta_{L_n})\end{aligned}$$



# Variational Symmetries

Symmetries arise from Lie group actions.

EXAMPLE:  $G = (\mathbb{R}, +)$

$$\epsilon \cdot x = x^* = \frac{x}{1 - \epsilon x}, \quad \epsilon \cdot u = u^*(x^*) = \frac{u(x)}{1 - \epsilon x}$$

Group Action Property

$$\begin{aligned} \delta \cdot (\epsilon \cdot x) &= \delta \cdot \left( \frac{x}{1 - \epsilon x} \right) = \frac{\frac{x}{1 - \epsilon x}}{1 - \delta \frac{x}{1 - \epsilon x}} \\ &= \frac{x}{1 - (\epsilon + \delta)x} = (\epsilon + \delta) \cdot x \end{aligned}$$

and similarly for  $u(x)$ .

Prolonged Group Action

$$\epsilon \cdot u_x = u_{x^*}^* = \frac{\partial u^*(x^*)}{\partial x} / \frac{\partial x^*}{\partial x} = \frac{u_x}{(1 - \epsilon x)^2}$$

and

$$\delta \cdot (\epsilon \cdot u_x) = \frac{\delta \cdot u_x}{(1 - \epsilon(\delta \cdot x))^2} = \frac{u_x}{(1 - (\delta + \epsilon)x)^2}$$

## Action on Integrals

$$\epsilon \cdot \int_{\Omega} L(x, u, u_x, \dots) dx$$

def'n of  $\epsilon \cdot$

$$= \int_{\epsilon \cdot \Omega} L(\epsilon \cdot x, \epsilon \cdot u, \epsilon \cdot u_x, \dots) d\epsilon \cdot x$$

change of variable

$$= \int_{\Omega} L(\epsilon \cdot x, \epsilon \cdot u, \epsilon \cdot u_x, \dots) \frac{d\epsilon \cdot x}{dx} dx$$

Use  $L^2$  theory to get that a variational symmetry of a Lagrangian is a group action such that

$$L(x, u, u_x, \dots) = L(\epsilon \cdot x, \epsilon \cdot u, \epsilon \cdot u_x, \dots) \frac{d\epsilon \cdot x}{dx}$$

## Infinitesimal Action on Integrals

Since the symmetry invariance condition

$$L(x, u, u_x, \dots) = L(\epsilon \cdot x, \epsilon \cdot u, \epsilon \cdot u_x, \dots) \frac{d\epsilon \cdot x}{d\epsilon}$$

is true all  $\epsilon$ , then if everything is sufficiently smooth, applying  $\frac{d}{d\epsilon}|_{\epsilon=0}$  to both sides, and noting that when  $\epsilon = 0$  we have the identity action,

$$\begin{aligned} 0 &= \frac{\partial L}{\partial x} \xi + \frac{\partial L}{\partial u} \phi + \frac{\partial L}{\partial u_x} \phi^x + \dots + L \xi_x \\ &= \frac{D(L\xi)}{Dx} + \frac{\partial L}{\partial u} Q + \frac{\partial L}{\partial u_x} \frac{DQ}{Dx} + \dots + \frac{\partial L}{\partial u_{xx}} \frac{D^2 Q}{Dx^2} + \dots \end{aligned}$$

where

$$Q = \phi - u_x \xi, \quad \phi = \frac{d}{d\epsilon}|_{\epsilon=0} \epsilon \cdot u, \quad \xi = \frac{d}{d\epsilon}|_{\epsilon=0} \epsilon \cdot x$$

and  $\frac{D}{Dx}$  is the total derivative operator.

$$0 = \text{Div}(L\xi) + \sum (D^J Q^\alpha) \frac{\partial L}{\partial u_J^\alpha}$$

Almost to the punchline

Let

$$\mathbf{v}_Q = \sum_{\alpha} Q^{\alpha} \frac{\partial}{\partial u^{\alpha}}$$

Then the *prolongation* is defined by

$$\text{pr}\mathbf{v}_Q = \sum_{\alpha, J} \mathbb{D}^J Q^{\alpha} \frac{\partial}{\partial u_J^{\alpha}}$$

Note

$$u_J^{\alpha} = \frac{\partial u^{\alpha}}{\partial x_1^{J_1} \cdots \partial x_p^{J_p}} = \mathbb{D}^J u^{\alpha}$$

Then

$$\sum \left( \mathbb{D}^J Q^{\alpha} \right) \frac{\partial L}{\partial u_J^{\alpha}} = \text{pr}\mathbf{v}_Q \lrcorner \hat{\mathbf{d}}L$$

Recall that  $\hat{\mathbf{d}}$  is one of the two operators comprising the Euler Lagrange operator, while the left hand side is a divergence if  $Q$  is the characteristic of a symmetry.

# THE PUNCHLINE

$$\begin{array}{c}
 \dots \xrightarrow{\text{Curl}} \wedge^2 \xrightarrow{\text{Div}} \wedge^3 \xrightarrow{E} \wedge_*^1 \xrightarrow{\pi \circ \hat{d}} \dots \\
 \qquad \qquad \qquad \Psi \qquad \qquad \qquad \Psi \\
 \qquad \qquad \qquad L \qquad \qquad \qquad E^\alpha(L) du^\alpha dx \\
 \qquad \qquad \qquad \qquad \qquad \qquad = \hat{d}(L) + \text{Div}(\eta_L) \\
 \qquad \qquad \qquad \overleftarrow{\text{pr}} \mathbf{v}_Q \lrcorner
 \end{array}$$

$$\begin{aligned}
 & Q \cdot E(L) \\
 &= \mathbf{v}_Q \lrcorner \hat{d}(L) + \text{Div}(\overleftarrow{\text{pr}} \mathbf{v}_Q \lrcorner \eta_L)
 \end{aligned}$$

If  $Q$  is the characteristic of a symmetry, we have that

$$\mathbf{v}_Q \lrcorner \hat{d}(L) = \text{Div}(L\xi)$$

and hence that

$$Q \cdot E(L) = \text{Div}(\text{something})$$

## Non-trivial example

### Semi-geostrophic equations

$$\begin{array}{ll}
 \text{Group} & \left\{ \begin{array}{l} a^* = \phi(a, b) \\ b^* = \psi(a, b) \end{array} \right. \quad \phi_a \psi_b - \phi_b \psi_a = 1 \\
 \text{Invariants} & \left\{ \begin{array}{l} h = (x_a y_b - x_b y_a)^{-1} \\ \partial_x = h(y_b \partial_a - y_a \partial_b) \\ \partial_y = h(-x_b \partial_a + x_a \partial_b) \end{array} \right. \\
 \text{Equations} & \left\{ \begin{array}{l} D_t x = -\frac{g}{f^2} D_t h_x - \frac{g}{f} h_y \\ D_t y = -\frac{g}{f^2} D_t h_y + \frac{g}{f} h_x \end{array} \right.
 \end{array}$$

The Lagrangian has 4 arbitrary functions which obey two conditions. The conserved quantity is *potential vorticity*

$$\frac{1}{h} \left( f + \frac{g}{f} (h_{xx} + h_{yy}) \frac{g^2}{f^3} (h_{xx} h_{yy} - h_{xy}^2) \right)$$

# DISCRETE Almost Punchline

This case is easier than the smooth case.

- Since  $n$  cannot vary in a smooth way, the “mesh variables”  $x_n$  are treated as dependent variables.
- The group action commutes with shift:

$$\epsilon \cdot S^j(u_n) = \epsilon \dot{u}_{n+j} = S^j \epsilon \cdot u_n$$

so no prolongation formulae are required.

For example,

$$\epsilon \cdot u_n = \frac{u_n}{1 - \epsilon x_n} \implies \epsilon \cdot u_{n+j} = \frac{u_{n+j}}{1 - \epsilon x_{n+j}}$$

The symmetry condition is:

$$\begin{aligned} L_n(x_n, \dots x_{n+j}, u_n, \dots u_{n+k}) \\ = L_n(x_n^*, \dots x_{n+j}^*, u_n^*, \dots u_{n+k}^*) \end{aligned}$$

where  $( )^* \equiv \epsilon \cdot ( )$ .

Applying

$$\frac{d}{d\epsilon}\bigg|_{\epsilon=0}$$

to both sides of the symmetry condition yields

$$0 = \sum_k \frac{\partial L_n}{\partial x_{n+k}} \frac{d}{d\epsilon}\bigg|_{\epsilon=0} x_{n+k}^* + \frac{\partial L_n}{\partial u_{n+k}} \frac{d}{d\epsilon}\bigg|_{\epsilon=0} u_{n+k}^*$$

Setting

$$Q_n^x = \frac{d}{d\epsilon}\bigg|_{\epsilon=0} x_n^*, \quad Q_n^u = \frac{d}{d\epsilon}\bigg|_{\epsilon=0} u_n^*$$

then since

$$Q_{n+k}^x = S^k(Q_n^x), \quad Q_{n+k}^u = S^k(Q_n^u)$$

the equation above can be written as

$$0 = X_Q \lrcorner \hat{d}L_n, \quad X_Q = \sum_{\alpha, J} S^J(Q_n^\alpha) \frac{\partial}{\partial u_{n+J}^\alpha}$$



## DISCRETE Punchline

$$\begin{array}{ccc}
 \xrightarrow{\Delta} \mathbf{Ex}^2 \xrightarrow{\Delta} \mathbf{Ex}^3 & \xrightarrow{\pi \circ \hat{d}} & \Lambda_*^1 \rightarrow \Lambda_*^2 \\
 \downarrow \Psi & & \downarrow \Psi \\
 & & E(L_n) du_n \\
 & & = \hat{d}(L_n) + \Delta(\eta_{L_n}) \\
 & \xleftarrow{X_{Q \lrcorner}} & 
 \end{array}$$

$$Q \cdot E(L_n) = X_{Q \lrcorner} \hat{d}(L_n) + \Delta(X_{Q \lrcorner} \eta_{L_n})$$

Again, we get that if

$$X_{Q \lrcorner} \hat{d}(L_n) = 0$$

then

$$Q \cdot E(L_n) = \Delta(\text{something}),$$

that is, a total difference expression which is zero on solutions of the discrete Euler Lagrange system.

**Nice example** T.D. Lee, Difference Equations and Conservation Laws, J. Stat. Phys., **46** (1987)

A difference model for  $\int (\frac{1}{2}\dot{x}^2 - V(x)) dt$

Define

$$\bar{V}(n) = \frac{1}{x_n - x_{n-1}} \int_{x_{n-1}}^{x_n} V(x) dx$$

and take

$$L_n = \left[ \frac{1}{2} \left( \frac{x_n - x_{n-1}}{t_n - t_{n-1}} \right)^2 - \bar{V}(n) \right] (t_n - t_{n-1})$$

The group action is  $t_n^* = t_n + \epsilon$ , with  $x_n$  invariant. The conserved quantity is thus “energy”. Now,  $Q_n^t = 1$  for all  $n$ , and  $Q_n^x = 0$ . The equations become

$$0 = E^t(L_n) = \frac{\partial}{\partial t_n} L_n + S \left( \frac{\partial}{\partial t_{n-1}} L_n \right)$$

$$0 = X_{Q^t} \lrcorner d(L_n) = \frac{\partial}{\partial t_n} L_n + \frac{\partial}{\partial t_{n-1}} L_n$$

as  $L_n$  is a function of  $(t_n - t_{n-1})$ .

It is easy to see in this case that

$$0 = (S - \text{id}) \left( \frac{\partial}{\partial t_n} L_n \right)$$

is implied by the two equations, to yield

$$\frac{1}{2} \left( \frac{x_n - x_{n-1}}{t_n - t_{n-1}} \right)^2 + \bar{V}(n) = c$$

Note that the energy in the smooth case is

$$1/2\dot{x}^2 + V.$$

Can regard the EL eqn for the mesh variables as an equation for a variable mesh.

## INTERLUDE

If we know the group action for a particular conservation law, we can “design in” that conservation law into a discretisation by taking a Lagrangian composed of invariants. These necessarily satisfy  $v_Q(I) = 0$  or  $X_Q(I_n) = 0$ . The Fels and Olver formulation of moving frames is particularly helpful here: a sample theorem is

### Discrete rotation invariants in $\mathbb{Z}^2$

Let  $(x_n, y_n), (x_m, y_m)$  be two points in the plane. Then

$$I_{n,m} = x_n y_n + x_m y_m, \quad J_{n,m} = x_n y_m - x_m y_n$$

are rotation invariants. Moreover, any discrete rotation invariant is a function of these.

## Made up example

Suppose

$$L_n = \frac{1}{2} J_{n,n+1}^2 = \frac{1}{2} (x_n y_{n+1} - x_{n+1} y_n)^2$$

then

$$\begin{cases} E_n^x &= J_{n,n+1} y_{n+1} - J_{n-1,n} y_{n-1} \\ E_n^y &= -J_{n,n+1} x_{n+1} + J_{n-1,n} x_{n-1} \end{cases}$$

Now,

$$Q_n = (Q_n^x, Q_n^y) = (-y_n, x_n) = \left. \frac{d}{d\theta} \right|_{\theta=0} (x_n^*, y_n^*)$$

and thus

$$\begin{aligned} Q_n \cdot E_n &= J_{n,n+1} (-y_n y_{n+1} - x_n x_{n+1}) \\ &\quad + J_{n-1,n} (y_n y_{n-1} + x_n x_{n-1}) \\ &= -J_{n,n+1} I_{n,n+1} + J_{n-1,n} I_{n-1,n} \\ &= -(S - \text{id})(J_{n-1,n} I_{n-1,n}) \end{aligned}$$

gives the conserved quantity.

Note that  $I_{n,m} = I_{m,n}$  and  $J_{n,m} = -J_{m,n}$

## Less easy example

Hereman et al., Densities, Symmetries and Recursion operators for nonlinear DDEs, CRM Proceedings

The Toda lattice in polynomial form is

$$\begin{cases} \dot{u}_n = v_{n-1} - v_n \\ \dot{v}_n = v_n(u_n - u_{n+1}) \end{cases}$$

The scaling symmetry is the basis for the ansatz used to obtain the differential-difference conservation laws, which are of the form

$$\frac{D}{Dt}\rho_n + (S - \text{id})J_n = 0$$

for example

$$\rho_n = \frac{1}{3}u_n^3 + u_n(v_{n-1} - v_n), J_n = u_{n-1}u_nv_{n-1} + v_{n-1}^2$$

These results use the ansatz plus homotopy operator method outlined earlier.

## Summary of the Pattern

$$\left\{ \begin{array}{c} \text{Div} \\ \Delta \end{array} \right\} \xrightarrow{\quad} \left\{ \begin{array}{c} \Lambda^3 \\ \mathbf{Ex}^3 \\ \Psi \\ L \end{array} \right\} \xrightarrow{\pi \circ \widehat{\mathbf{d}}} \Lambda_*^1 \longrightarrow \Lambda_*^2 \longrightarrow \Psi \sum_{\alpha} E^{\alpha}(L) \mathrm{d} u^{\alpha} = \widehat{\mathbf{d}}(L) + \left\{ \begin{array}{c} \text{Div} \\ \Delta \end{array} \right\} \eta_L$$

$$\left\{ \begin{array}{c} \leftarrow \\ v_Q \\ X_Q \end{array} \right\}_{\perp}$$

$$Q \cdot E(L) = \begin{Bmatrix} v_Q \\ X_Q \end{Bmatrix} \lrcorner \hat{\mathbf{d}}L + \begin{Bmatrix} \text{Div} \\ \Delta \end{Bmatrix} \begin{Bmatrix} v_Q \\ X_Q \end{Bmatrix} \lrcorner \boldsymbol{\eta}_L$$

- the formula for  $\eta_L$  is explicit, exact, symbolic
- the first summand is a total derivative or difference by the symmetry condition

# OK let's try for a Neother's Theorem for Finite Element!

D. Arnold, Beijing ICM Plenary talk

Given a system of moments and sundry other data, aka degrees of freedom, that yield projection operators such that the diagram commutes:

$$\begin{array}{ccccccccc} 0 & \rightarrow & \mathbb{R} & \rightarrow & \Lambda^0 & \rightarrow & \Lambda^1 & \rightarrow & \Lambda^2 & \rightarrow & \Lambda^3 & \rightarrow & 0 \\ & & & & \Pi_0 \downarrow & & \Pi_1 \downarrow & & \Pi_2 \downarrow & & \Pi_3 \downarrow & & \\ 0 & \rightarrow & \mathbb{R} & \rightarrow & \mathcal{F}^0 & \rightarrow & \mathcal{F}^1 & \rightarrow & \mathcal{F}^2 & \rightarrow & \mathcal{F}^3 & \rightarrow & 0 \end{array}$$

all relative to some triangulation.

**Yields stability!!** A Lagrangian is composed of wedge products of 1-, 2- and 3- forms.

Choose the discretisation of each to be in the relevant  $\mathcal{F}_i$ . Then commutativity implies conditions for Brezzi's theorem to hold.



In one dimension: with  $e_n = (x_n, x_{n+1})$ ,  $\Pi_0$  to piecewise linear,  $\Pi_1$  to piecewise constant with moment

$$\alpha_n = \int_{x_n}^{x_{n+1}} u(x) \psi_n(x) \, dx$$

Commutativity of the diagram

$$\begin{array}{ccc} u & \xrightarrow{d} & u_x dx \\ \Pi_0 \downarrow & & \downarrow \Pi_1 \\ u|_{e_n} = A_n x + B_n & \mapsto & A_n = \int_{x_n}^{x_{n+1}} u'(x) \psi_n(x) \, dx \end{array}$$

implies

$$A_n = [u(x) \psi_n(x)]_{x_n}^{x_{n+1}} - \int_{x_n}^{x_{n+1}} u(x) \psi_n'(x) \, dx$$

Note that

$$\int_{x_n}^{x_{n+1}} \psi_n(x) \, dx = 1.$$

is required by the projection property.

A finite element Lagrangian is built up of wedge products of forms in  $\mathcal{F}_0, \mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$ . Call this resulting space  $\tilde{\mathcal{F}}_3$ . In each top-dimensional simplex, denoted  $\tau$ , integrate to get

$$L = \sum_{\tau} L_{\tau}(\alpha_{\tau}^1, \dots, \alpha_{\tau}^p)$$


where  $\alpha_{\tau}^j$  is the  $j^{\text{th}}$  degree of freedom in  $\tau$ .  $L$  can also depend on mesh data.

Can now take  $\hat{d}$  which is the variation with respect to the  $\alpha_{\tau}^j$ .

**EXAMPLE** In one dimension,

$$\begin{array}{ccccccc}
 0 & \rightarrow & \mathbb{R} & \rightarrow & \Lambda^0 & \xrightarrow{d} & \Lambda^1 \rightarrow 0 \\
 & & & & \Pi_0 \downarrow & & \Pi_1 \downarrow \\
 0 & \rightarrow & \mathbb{R} & \rightarrow & \mathcal{F}_0 & \xrightarrow{d} & \mathcal{F}_1 \rightarrow 0
 \end{array}$$

$\Pi_1$  is to piecewise constant functions with moment  $\bar{u}(n) = \int_{x_n}^{x_{n+2}} u(x) \psi_n(x) dx$  where



on  $(x_n, x_{n+2})$ , while  $\Pi_0$  is to piecewise linear functions with moments

$$\alpha_n = \frac{1}{x_n - x_{n+2}} \int_{x_n}^{x_{n+1}} u(x) dx$$

that is,  $\alpha_n, \alpha_{n+1}$  are used in  $(x_n, x_{n+2})$ ;

$$\begin{aligned}
 u \mapsto & 2 \frac{\alpha_{n+1} - \alpha_n}{x_{n+2} - x_n} x + \left( \frac{x_{n+1} + x_{n+2}}{x_{n+2} - x_n} \right) \alpha_n \\
 & - \left( \frac{x_{n+1} + x_n}{x_{n+2} - x_n} \right) \alpha_{n+1}
 \end{aligned}$$

## Very simple example

$\int \frac{1}{2} u_x^2 dx$  projects to

$$\sum_n \int_{x_{2n}}^{x_{2n+2}} \frac{1}{2} \Pi(u)_x^2 dx = \sum_n 2 \left( \frac{(\alpha_{2n} - \alpha_{2n+1})^2}{x_{2n+2} - x_{2n}} \right)$$

Then

$$\begin{aligned} \hat{d}L_{2n} &= 4 \frac{\alpha_{2n} - \alpha_{2n+1}}{x_{2n+2} - x_{2n}} (d\alpha_{2n} - d\alpha_{2n+1}) \\ &= 4 \left( \frac{\alpha_{2n} - \alpha_{2n+1}}{x_{2n+2} - x_{2n}} - \frac{\alpha_{2n+1} - \alpha_{2n+2}}{x_{2n+2} - x_{2n+1}} \right) d\alpha_{2n} \\ &\quad + (S - \text{id})(\text{something}) \end{aligned}$$

The discrete Euler Lagrange equation is then, after “integration”,

$$\frac{\alpha_{2n} - \alpha_{2n+1}}{x_{2n+2} - x_{2n}} = c$$

Look now at the “Noether pattern” for the Finite Element variational complex

$$\begin{array}{ccccccc}
 \rightarrow \tilde{\mathcal{F}}_2 & \rightarrow & \tilde{\mathcal{F}}_3 & \xrightarrow{\pi \circ \hat{d} \circ \int} & \tilde{\mathcal{F}}_*^1 & \rightarrow & \tilde{\mathcal{F}}_*^2 \rightarrow \\
 & & \downarrow \Psi & & \downarrow \Psi & & \\
 & & L_\tau & & E(L_\tau) + \delta(\eta_L) & & \\
 & & & & = \hat{d}L_\tau & & \\
 & & & & \downarrow v_{Q_\tau \lrcorner} & & 
 \end{array}$$

where  $\delta$  is the mesh dependent coboundary operator (recall  $\delta(f)(\tau) = f(\partial\tau)$ ).

Step 1: find  $\eta_L$

Step 2: find  $v_Q$

If then  $v_{Q_\tau \lrcorner} \hat{d}(L_\tau) = \delta(\text{something})$  we will have that

$$0 = Q_\tau \cdot E(L_\tau) + \delta(\text{something}).$$

## Group actions on moments

The clue is the variational symmetry group action on  $\int_{\Omega} L(x, u, \dots) dx$

Define

$$\begin{aligned}\epsilon \cdot \int_{\tau} u(x) \psi_{\tau}(x) dx \\ = \int_{\tau} \epsilon \cdot u(x) \psi_{\tau}(\epsilon \cdot x) \frac{d\epsilon \cdot x}{dx} dx\end{aligned}$$

Example Recall the projective action

$$\epsilon \cdot x = \frac{x}{1 - \epsilon x}, \quad \epsilon \cdot u(x) = \frac{u(x)}{1 - \epsilon x}$$

Then the induced action on the moments

$$\alpha_n = \int_{x_n}^{x_{n+1}} \frac{u(x)}{x^3} dx, \quad \beta_n = \int_{x_n}^{x_{n+1}} \frac{u(x)}{x^4} dx$$

is

$$\epsilon \cdot \alpha_n = \alpha_n, \quad \epsilon \cdot \beta_n = \beta_n - \epsilon \alpha_n$$

In general for this action,

$$\begin{aligned} & \epsilon \cdot \int_{x_n}^{x_{n+1}} x^m u(x) \, dx \\ &= \int_{x_n}^{x_{n+1}} \frac{x^m}{(1-\epsilon x)^m} \frac{u(x)}{1-\epsilon x} \frac{dx}{(1-\epsilon x)^2} \\ &= \int_{x_n}^{x_{n+1}} \frac{x^m u(x)}{(1-\epsilon x)^{m+3}} \, dx \end{aligned}$$

THINK: if you want a coherent scheme which maps to itself under this projective action, and involves only a finite amount of data, then take your moments to be

$$u(x) \mapsto \int_{x_n}^{x_{n+1}} \frac{u(x)}{x^m} \, dx, \quad m = 3, 4, \dots, N.$$

## CONCLUSIONS

- The underlying algebraic pattern of the exact variational complexes provide a framework for generalisations of Noether's Theorem and conservation laws in general.
- Symmetry-adapted moments would appear to be necessary.
- Next: formulae for  $\eta_{L_\tau}$  where

$$\hat{d}(L_\tau) = E(L_\tau) + \delta(\eta_{L_\tau})$$

in terms of the mesh dependent coboundary operator.